

# Dietary Supplements for Health, Adaptation, and Recovery in Athletes

**Eric S. Rawson**  
Messiah College

**Mary P. Miles**  
Montana State University

**D. Enette Larson-Meyer**  
University of Wyoming

Some dietary supplements are recommended to athletes based on data that supports improved exercise performance. Other dietary supplements are not ergogenic per se, but may improve health, adaptation to exercise, or recovery from injury, and so could help athletes to train and/or compete more effectively. In this review, we describe several dietary supplements that may improve health, exercise adaptation, or recovery. Creatine monohydrate may improve recovery from and adaptation to intense training, recovery from periods of injury with extreme inactivity, cognitive processing, and reduce severity of or enhance recovery from mild traumatic brain injury (mTBI). Omega 3-fatty acid supplementation may also reduce severity of or enhance recovery from mTBI. Replenishment of vitamin D insufficiency or deficiency will likely improve some aspects of immune, bone, and muscle health. Probiotic supplementation can reduce the incidence, duration, and severity of upper respiratory tract infection, which may indirectly improve training or competitive performance. Preliminary data show that gelatin and/or collagen may improve connective tissue health. Some anti-inflammatory supplements, such as curcumin or tart cherry juice, may reduce inflammation and possibly delayed onset muscle soreness (DOMS). Beta-hydroxy beta-methylbutyrate (HMB) does not consistently increase strength and/or lean mass or reduce markers of muscle damage, but more research on recovery from injury that includes periods of extreme inactivity is needed. Several dietary supplements, including creatine monohydrate, omega 3-fatty acids, vitamin D, probiotics, gelatin, and curcumin/tart cherry juice could help athletes train and/or compete more effectively.

**Keywords:** creatine, curcumin, gelatin, HMB, omega 3 fatty acids, probiotics, tart cherry juice, vitamin D

Several dietary supplements, including carbohydrate, caffeine, creatine monohydrate, nitrate, beta-alanine, and sodium bicarbonate, are well-established performance enhancers (see [Peeling et al., 2018](#)). Additionally, the beneficial effects of protein on the adaptive response to exercise has been well studied (see [Hector & Phillips, 2018](#)). Further, some supplements, such as caffeine and carbohydrate, may augment both muscle and brain performance (see [Peeling et al., 2018](#)). There is also research indicating that some other dietary supplements may be valuable in enhancing muscular adaptations to exercise, improving brain performance, decreasing delayed onset muscle soreness or pain, reducing injury severity, enhancing recovery from injury, reducing gastrointestinal problems, and decreasing respiratory tract infection illness load. For the most part, these effects are not ergogenic, but may help athletes to train and/or compete more effectively without performance impediments. These supplements include creatine monohydrate,

beta-hydroxy beta-methylbutyrate (HMB), omega-3 fatty acids, vitamin D, probiotics, gelatin, and anti-inflammatory supplements such as curcumin or tart cherry juice. The potential benefits of these dietary supplements are summarized in Table 1.

## Creatine Monohydrate

### Muscular Adaptations

The ingestion of the dietary supplement creatine monohydrate (about 20 g/day for 5 days) increases muscle creatine by about 20% ([Harris et al., 1992](#); [Hultman et al., 1996](#)). Subsequently, the performance of sports and exercises that rely heavily on creatine and phosphorylcreatine to resynthesize adenosine triphosphate (ATP) (e.g., <30 s, intense, repeated bouts) can be improved (reviewed in [Branch, 2003](#); [Gualano et al., 2012](#); [Rawson & Volek, 2003](#)). In addition to the well-established performance-enhancing effects, creatine monohydrate also works through multiple mechanisms to enhance muscle recovery from intense exercise (reviewed in [Heaton et al., 2017](#)).

Muscle phosphorylcreatine and glycogen can be depleted following intense exercise, but creatine supplementation can enhance replenishment of these fuel sources. Faster postexercise

---

Rawson is with the Department of Health, Nutrition, and Exercise Science, Messiah College, Mechanicsburg, PA. Miles is with the Department of Health and Human Development, Montana State University, Bozeman, MT. Larson-Meyer is with the Nutrition & Exercise Laboratory, Department of Family and Consumer Sciences, University of Wyoming, Laramie, WY. Address author correspondence to Eric S. Rawson at [erawson@messiah.edu](mailto:erawson@messiah.edu).

**Table 1 Dietary Supplements for Health, Adaptation, and Recovery in Athletes**

Supplement/Nutrient	Deficiency	Dysfunction With Deficiency	Potential Benefits of Supplementation	Comments
Creatine monohydrate: Creatine is a naturally occurring nutrient, consumed in the diet, and synthesized in the body. Recommended supplement dose is 20 g/day for 5 days followed by 3 to 5 g/day to increase and maintain elevated body creatine levels (Harris et al., 1992; Hultman et al., 1996).	Low dietary creatine intake (e.g., vegan, vegetarian, or low meat eater) decreases muscle and blood muscle creatine concentrations.	No adverse clinical or performance findings with dietary-induced deficiency.	Increased lean mass/strength; enhanced recovery from intense exercise (faster PCr and glycogen resynthesis); enhanced adaptive response to exercise via increased growth factor/gene expression, increased intracellular water; reduced symptoms of or enhanced recovery from muscle damaging exercise (e.g., DOMS); enhanced recovery from disuse, immobilization, or extreme inactivity such as after injury; improved cognitive processing; decreased risk/enhanced recovery from mTBI.	Well studied. Excellent safety profile. A small increase in body mass is common with supplementation. This may be relevant for sports with weight classes/restrictions or where increased body mass may decrease performance. Not known if improved cognitive processing translates into improved athletic performance.
Beta-hydroxy Beta-methylbutyrate (HMB): HMB is a metabolite of the amino acid leucine. Manufacturer recommended dosage is 3 g/day (Rowlands & Thomson, 2009; Sanchez-Martinez et al., 2017).	N/A/	N/A	Enhanced lean mass, strength, and adaptation to exercise via decreased muscle protein breakdown; reduced symptoms of or enhanced recovery from muscle damaging exercise (e.g., decreased DOMS).	Cannot be confidently recommended to athletes. Effects may be no more effective than adhering to the current protein intake recommendations. May prove useful in rehabilitation where there may be periods of extreme inactivity, but few data are available.
Vitamin D: An essential fat-soluble vitamin. Skin exposure to sunlight accounts for 90% of the source of vitamin D.	Insufficiency/deficiency has been identified in athletes from recreational through professional levels especially, but not exclusively, when access to sunlight is limited	Blunted adapted to response to exercise likely; increased URTI; increased injury, stress fractures.	Improved adaptive response to exercise; decreased incidence, duration, and severity of URTI; decreased stress fractures.	RDA 600 IU/d; identify deficiencies through blood work, consult with a physician and sport nutritionist, and take corrective actions in insufficient/deficient athletes with changes in lifestyle, food intake, and possibly supplementation; indiscriminate vitamin D supplementation for athletes is not advised; compensatory response to high-dose supplementation possible (e.g., decreased PTH), but not well studied.
Omega 3-fatty acids: Dose most commonly studied is about 2 g/day (Jeromson et al., 2015; Mickleborough, 2013).	Not described in athletes.	Not described in athletes.	Improved cognitive processing; decreased risk/enhanced recovery from mTBI; reduced symptoms of or enhanced recovery from muscle damaging exercise (e.g., DOMS).	Not known if improved cognitive processing translates into improved athletic performance; few data on mTBI, but animal data and two case studies show that the structural damage and cognitive decline associated with mTBI are reduced/attenuated with omega-3 fatty acid; benefits to muscle damage are not a consistent finding; low risk but unclear if supplementation should be pursued by athletes, in lieu of including fatty fish in the diet as a source of omega-3 fatty acids.
Probiotics: Effective dosing regimens of $10^9$ to $4 \times 10^{10}$ CFU for 4 to 21 weeks in duration (Pyne et al., 2015).	N/A	N/A	Decreased severity or duration of gastrointestinal problems; decreased incidence, duration, and severity of URTI.	Modest benefits to athletes prone to GIT problems or traveling to regions in which GIT problems are more likely; most studies report reduced incidence, duration, and severity of URTI in athlete; many different probiotic supplements have been studied, so specific recommendations and conclusions are difficult to ascertain; supplementation needs to begin well ahead of competition.

(continued)

Supplement/Nutrient	Deficiency	Dysfunction With Deficiency	Potential Benefits of Supplementation	Comments
Gelatin and vitamin C/ collagen: Recommended dose of gelatin is 5 to 15 g gelatin w/50 mg vitamin C (Shaw et al., 2017). Collagen hydrolysate dose is about 10 g/day (Clark et al., 2008; McAlindon et al., 2011).	N/A	N/A	Increased collagen production, thickened cartilage, decreased knee pain.	Gelatin and collagen supplements appear to be low risk; few data are available; functional benefits, recovery from injury, and effects in elite athletes are not known.
Anti-inflammatory supplements: Curcumin is a constituent of the spice turmeric, and supplements are often ingested at a dose of about 5 g/day (McFarlin et al., 2016; Nicol et al., 2015). Tart cherry juice is ingested at a dose of about 8 to 12 oz. (1 oz. if concentrate) twice/day for 4 to 5 days before an athletic event or for 2 to 3 days afterwards to promote recovery (Bell et al., 2014, Coelho Rabello Lima et al., 2015).	N/A	N/A	Anti-inflammatory effects; reduced symptoms of or enhanced recovery from muscle damaging exercise (e.g., decreased DOMS).	Reduced DOMS may be important in sports where soreness might impair performance in a subsequent bout of exercise; benefits may be sport/training specific; more research needs to be conducted before these compounds could be recommended to athletes.

phosphorylcreatine resynthesis may improve recovery from an acute bout of sprint type exercise, or during repeated bouts of sprints, which mimics the stop and go, intermittent nature of many sports. Consequently, this could improve exercise performance in subsequent bout(s) a few minutes later (Greenhaff et al., 1994; Yquel et al., 2002). Also, simultaneous creatine and carbohydrate loading further increases glycogen resynthesis (12%) (Nelson et al., 2001), with 82% of this increase occurring during the first 24 hr (Roberts et al., 2016) (reviewed in Volek & Rawson, 2004). This could enhance the performance of a subsequent bout of exercise occurring hours or days later.

cell survival (Safdar et al., 2008). Some of these effects may be modulated by increased intramuscular water (Deminice et al., 2016), which can inhibit protein breakdown and RNA degradation, and stimulate protein, DNA, and RNA synthesis (Berneis et al., 1999; Häussinger et al., 1993).

## Recovery from Disuse/Immobilization

Along with decreases in muscle mass and function, muscle creatine decreases as much as 24% during periods of extreme inactivity, such as during immobilization (MacDougall, Ward, Sale, & Sutton, 1977). Consequently, maintaining or increasing muscle creatine with supplementation during periods of inactivity or recovery from injury may prove advantageous. Seven studies of the effects of creatine supplementation during immobilization have been

### $\beta$ -Hydroxy $\beta$ -Methylbutyrate (HMB)

HMB is ingested in the diet and is a metabolite of leucine. Animal research indicates that HMB can increase body mass in slaughter animals (reviewed in [Szczesniak et al., 2015](#)). As a supplement for humans, HMB is marketed as an anticatabolic nutrient that decreases protein breakdown; a dose of about 3 g/day has been recommended. Potential mechanisms underlying anticatabolic effects include decreased protein breakdown through reduced expression of components of the ubiquitin proteasome pathway, increased protein synthesis through phosphorylation of mammalian target rapamycin (mTOR) substrates and phosphorylation of eukaryotic initiation factor (eIF2), increased cholesterol synthesis, increased growth hormone and IGF I mRNA, increased proliferation and differentiation of satellite cells, and inhibited apoptosis (reviewed in [Szczesniak et al., 2015](#)). However, these potential benefits may be just as easily obtained with ingestion of leucine or whey protein, which increase muscle protein synthesis and anabolic signaling ([Wilkinson et al., 2013](#)).

If HMB is anticatabolic, it could improve training adaptations by decreasing muscle damage or protein breakdown. The effects of HMB supplementation on markers of muscle damage and resistance training adaptations in lean mass and strength have been well reviewed (Fitschen, Wilson, Wilson, & Wilund, 2013; Molfino, Gioia, Rossi Fanelli, & Muscaritoli, 2013; Zanchi et al., 2011). While early research and a meta-analysis concluded that HMB supplementation both increased lean mass and decreased muscle damage (Nissen & Sharp, 2003; Nissen et al., 1996), these conclusions have not stood the test of time. In two meta-analyses, it was concluded that HMB supplementation had inconsequential or trivial effects on strength and fat free mass, especially in trained competitive athletes, and that effects on muscle damage (based on blood creatine kinase levels) were unclear (Rowlands & Thomson, 2009; Sanchez-Martinez et al., 2017). Unfortunately, data on HMB supplementation are discrepant and difficult to interpret due to clustering of data sources (i.e., positive studies from the same/similar labs) and reliance on nonspecific markers of muscle damage (e.g., creatine kinase) (Decombaz et al., 2003). Additionally, in some instances, potential conflicts of interest were not fully disclosed (Editor, 2003).

The recent publication of several studies on HMB-free acid (HMB-FA) supplementation where implausible anabolic drug-like gains in strength, power, and fat free mass, and reductions in muscle damage were reported (Lowery et al., 2016; Wilson et al., 2013; Wilson et al., 2014) has further complicated interpretation of the HMB literature. These findings have been challenged by other researchers (Gentles & Phillips, 2017; Phillips et al., 2017), but, at this time, the authors have not responded. HMB appears to have some biologic activity in humans that may aid in muscular adaptations from exercise, but it is doubtful that effects are as powerful as some studies have shown. Perhaps, HMB (reviewed in Wu et al., 2015), like creatine monohydrate (reviewed in Gualano et al., 2012; Gualano et al., 2016), is of benefit for patient populations or sarcopenic older adults. Unlike creatine monohydrate, HMB cannot be confidently recommended to athletes, as the effects may not be more effective than adhering to the current protein intake recommendations. HMB may prove useful in rehabilitation where there may be periods of extreme inactivity, as Deutz and others (2013) showed better preservation of lean mass in older adults ingesting HMB during 10 days of bed rest.

IJSNEM Vol. 28, No. 2, 2018

ultraviolet B (UVB) rays in sunlight (Hossein-nezhad & Holick, 2013). Synthesis of vitamin D in skin cells, however, is dependent on factors including time of day of exposure, season, latitude, cloud cover, smog, skin pigmentation, skin area exposed, sunscreen coverage, and age. Cutaneous vitamin D is not synthesized during the winter at latitudes greater than  $\sim 33^\circ$  north or south because insufficient UVB photons reach the earth's surface during these months (Hossein-nezhad & Holick, 2013).

Although there is disagreement over appropriate cut points of serum concentrations of 25-hydroxyvitamin D (the marker of vitamin D status), commonly used cut points define deficient as a serum 25 hydroxyvitamin D of <50 nmol/L, insufficient as <75 nmol/L, sufficient as >75 nmol/L, and optimal as 100 to 250 nmol/L (Larson-Meyer, 2015). Using these or similar cut points, it appears that many athletes are insufficient or deficient at various times of the year (Halliday et al., 2011; Larson-Meyer, 2015), including those who live in sunny environments close to the equator (Hamilton et al., 2010). Some data on elite and professional athletes also show a high percentage of deficiency and insufficiency (Allison, Close et al., 2015; Allison, Farooq et al., 2015; Close, Leckey et al., 2013; Close, Russell et al., 2013; Fishman et al., 2016; Maroon et al., 2015; Morton et al., 2012; Owens et al., 2017).

The data on the effects of vitamin D supplementation on muscle function have been described as equivocal, with discrepancies in research in part caused by differences in baseline 25-hydroxyvitamin D concentrations prior to supplementation (Close et al., 2016). For example, Close, Russell, et al. (2013) showed improved sprint time and vertical jump height following vitamin D supplementation in deficient athletes, but a separate study showed no effect of supplementation on strength (Owens et al., 2014). Owens and colleagues (2015) recently demonstrated that vitamin D supplementation (6 weeks; 4,000 IU/day) improved recovery of posteccentric exercise peak torque in vitamin D insufficient men and increased in vitro myotube hypertrophy. These data strongly suggest a role for adequate vitamin D in the adaptive process to intense exercise. Whether vitamin D supplementation is necessary for adequate adaptive effects likely depends on the athletes' vitamin D status at the initiation of supplementation.

In addition to muscle function, vitamin D supplementation may also be important for reducing risk of upper respiratory tract infections and stress fractures. These factors compromise an athlete's ability to train and compete. Studies in athletes (Cox et al., 2008; He, Aw Yong, Walsh, & Gleeson, 2016), military personnel (Laaksi et al., 2007), and the general population (Berry et al., 2011; Ginde et al., 2009; Sabetta et al., 2010) have reported negative associations between vitamin D status and incidences of upper respiratory tract illness (URTI). One study in college athletes found that serum 25(OH)D concentrations over the winter and spring were negatively associated with documented frequency of acute URTI (Halliday et al., 2011). The breakpoint for contracting a single episode occurred at ~95 nmol/l such that all athletes with concentrations lower than this breakpoint experienced one or more episodes of illness, whereas those with higher concentrations had one or fewer episodes. A similar study in endurance athletes reported that a greater proportion of athletes who maintained serum 25(OH)D concentrations of <30 nmol/L presented with URTI symptoms over this time point. The fewest symptoms were reported in those with 25(OH)D concentrations >120 nmol/L (He, Handzlik, et al., 2013). Athletes with low vitamin D concentrations also experienced a higher number of days with upper respiratory symptoms and higher symptom-severity scores. Although additional studies are needed to confirm the effectiveness of

additional studies are needed to confirm the effectiveness of

IJSNEM Vol. 28, No. 2, 2018

typically fall within the range of  $10^9$  to  $4 \times 10^{10}$  CFU for 4 to 21 weeks in duration (Pyne et al., 2015). The convention for naming bacteria is *Species strain* (e.g., *Lactobacillus rhamnosus* [*L. rhamnosus*]). A variety of strains within the *Lactobacillus*, *Bifidobacterium*, and, less frequently, the *Streptococcus* species are used as probiotics. These bacteria are Gram-positive (do not contain endotoxin).

Strenuous and prolonged exercise place stresses on the GIT that increase the likelihood of discomfort, abdominal cramping, acid reflux (heartburn), nausea, vomiting, diarrhea, and leakiness of the gut that may allow endotoxemia to occur (de Oliveira et al., 2014). Splanchnic hypoperfusion leading to ischemia in the gut is accepted as a principle cause, with additional contributions from nutritional, mechanical (e.g., jarring), and genetic influences that make some individuals more susceptible than others (de Oliveira et al., 2014). Probiotic support to increase resilience of the GIT against ischemia is of interest to athletes, particularly for those in prolonged endurance events that have the greatest occurrence of GIT problems that impair or stop performance. The proposed mechanism by which probiotics prevent GIT problems stemming from ischemia are related to improvement in the mucosal and epithelial barriers preventing a 'leaky gut' and in production of anti-inflammatory mediators. Probiotics do not eliminate hypoperfusion of the GIT, thus moderate reduction in severity and duration of gastrointestinal symptoms and exercise-induced endotoxemia are the best outcomes that might be expected in clinical trials. To this end, a number of clinical trials have measured small to moderate decreases in severity or duration of gastrointestinal problems attributed to use of *L. rhamnosus*, *L. fermentum*, and two different multi-strain probiotics (Haywood et al., 2014; Kekkonen et al., 2007; Shing et al., 2014; West et al., 2011). However, in the *L. fermentum* trial, benefits were lower in female participants, and the decrease in severity was accompanied by a two-fold greater increase in the number of and duration of less-severe symptoms in male participants for the probiotic compared to placebo group. The latter was proposed to be part of the adaptive response to probiotic colonization. Using a biomarker of gut permeability, a supporting clinical trial demonstrated that multi-strain probiotic reduced leakiness of the GIT as an underlying cause of GIT distress (Lamprecht et al., 2012). Upon reviewing the evidence, Pyne et al. (2015) reached the reasonable conclusion that athletes prone to GIT problems, or traveling to regions in which GIT problems are more likely, may benefit modestly from the above probiotics, but supplementation needs to begin well ahead of competitions and more research is needed before probiotics are recommended with confidence.

Microbes in the gut influence gastrointestinal tract (GIT) functions, the immune system, and metabolite production in a variety of ways that exert influence on exercise. The GIT microbiome primarily consists of a diverse array of bacterial species, each of which interacts with the host in a unique way. This influence ranges from beneficial, to neutral, to pathogenic or opportunistic, and the net effect of the GIT microbiome is a function of the combined effects of many different strains and their relative abundance. By definition, probiotics are strains of bacteria that can maintain viability in supplement form for oral ingestion, survive the acidic gastric environment, adhere and colonize in the GIT, antagonize pathogenic bacteria, and provide health benefit(s) to the host (athlete) (West et al., 2009). There are dozens of bacterial strains used as probiotics, particularly those that produce lactic acid, but each strain is unique with respect to how it responds to and affects the GIT milieu, the GIT, the immune system components within the GIT, and all downstream systems of the body. Thus, it is important to be specific about effects of each specific probiotic strain or combination of strains and not to generalize effects of one probiotic to all probiotics. Research regarding athletes has been undertaken to evaluate the efficacy of probiotic supplementation to (a) prevent or relieve GIT problems during exercise, (b) reduce the impact of URTIs, and (c) improve endurance performance.

While there is a variety of bacteria that have been used as probiotics in clinical studies in athletes, there is more consistency in the dosage and duration generally needed to produce benefits. A single bacterium is a colony-forming unit (CFU) and dosages are expressed using this common terminology. Dosing regimens

## Probiotics

Microbes in the gut influence gastrointestinal tract (GIT) functions, the immune system, and metabolite production in a variety of ways that exert influence on exercise. The GIT microbiome primarily consists of a diverse array of bacterial species, each of which interacts with the host in a unique way. This influence ranges from beneficial, to neutral, to pathogenic or opportunistic, and the net effect of the GIT microbiome is a function of the combined effects of many different strains and their relative abundance. By definition, probiotics are strains of bacteria that can maintain viability in supplement form for oral ingestion, survive the acidic gastric environment, adhere and colonize in the GIT, antagonize pathogenic bacteria, and provide health benefit(s) to the host (athlete) (West et al., 2009). There are dozens of bacterial strains used as probiotics, particularly those that produce lactic acid, but each strain is unique with respect to how it responds to and affects the GIT milieu, the GIT, the immune system components within the GIT, and all downstream systems of the body. Thus, it is important to be specific about effects of each specific probiotic strain or combination of strains and not to generalize effects of one probiotic to all probiotics. Research regarding athletes has been undertaken to evaluate the efficacy of probiotic supplementation to (a) prevent or relieve GIT problems during exercise, (b) reduce the impact of URTIs, and (c) improve endurance performance.

resistance, high-force eccentric, running, and cycling exercise (reviewed in [Bell et al., 2014](#); [Coelho Rabello Lima et al., 2015](#)). There are concerns that blunting inflammation may impair the adaptive response to exercise training. However, at times, such as when there are multiple competitions in a brief period of time, faster recovery of strength or decreased soreness may be more valued by the athlete than adaptation ([Bell et al., 2014](#)). Thus, it appears that curcumin and tart cherry juice may offer some benefits to athletes, although these benefits may be sport/training specific, and more research needs to be conducted before they could be recommended to athletes.

## Summary

In this review, several dietary supplements are described that can potentially play a role in health, exercise adaptation, and/or recovery for athletes. As summarized in Table 1, creatine monohydrate, vitamin D, omega 3-fatty acids, probiotics, gelatin/collagen, and certain anti-inflammatory supplements can influence cellular and tissue health, resilience, and repair in ways that may help athletes maintain health, adapt to exercise, and increase the quality and quantity of their training. Creatine monohydrate could be recommended for promoting recovery from, and muscular adaptations to, intense training; recovery from periods of injury that result in extreme inactivity; cognitive processing; and reducing severity of or enhancing recovery from mTBI. Vitamin D insufficiency or deficiency appears common, and the benefits of replenishment with supplementation likely improve some aspects of muscle function. However, athletes should seek a blood test and guidance from a physician to confirm inadequate vitamin D status, and then guidance from a nutrition professional regarding supplementation. There are several potential benefits of Omega fat supplementation on skeletal muscle, but more research needs to be conducted. Like creatine monohydrate, omega 3-fatty acid may reduce severity of or enhance recovery from mTBI. At this time, a recommendation for fatty fish consumption, which would include increased intake of omega 3-fatty acids, protein, vitamins, and minerals, may be more appropriate for athletes than a recommendation for fish oil supplementation. The majority of placebo-controlled clinical trials show that probiotic supplementation can reduce the incidence, duration, and severity of URTI in athletes. It is difficult to make more specific recommendations, as many different probiotic supplements were used in these studies. However, it is likely that there are several probiotic supplements that can improve training or competitive performance by decreasing the negative effects caused by URTI and/or GIT problems. A small amount of literature exists that demonstrates a benefit of gelatin and/or collagen on connective tissue health. As adverse effects from gelatin/collagen supplements appear low, at worst, these supplements are just an inexpensive source of amino acids, so the benefits of supplementation outweigh the risks. Anti-inflammatory supplements, such as curcumin and tart cherry juice, show promise in their ability to reduce markers of muscle damage and/or inflammation. However, it is unknown if these effects can enhance recovery in elite athletes or affect subsequent performance. HMB cannot be recommended at this time, but more research on recovery of muscle injury during periods of extreme inactivity is needed.

Much of the research on nutrition and the adaptive response to exercise focuses on dietary supplements that affect skeletal muscle tissue, MPS, and MPB. However, nutritional interventions that could benefit connective tissue proteins would be valuable as well (reviewed in [Baar, 2017](#)). Shaw et al. (2017) showed the combination of exercise plus a gelatin/vitamin C supplement (15 g gelatin plus 50 mg vitamin C) increased collagen production in vitro, and amino terminal propeptide (PINP) in the blood. Collagen hydrolysate supplementation (about 10 g/day) has been shown to thicken cartilage in osteoarthritis patients ([McAlindon et al., 2011](#)), and decrease knee pain in athletes ([Clark et al., 2008](#)). These studies suggest a benefit of gelatin plus vitamin C and/or collagen supplementation, but data on functional benefits in elite athletes, such as performance changes or recovery from injury, are not available.

A constituent of the spice turmeric, curcumin supplements are often ingested for anti-inflammatory effects at a dose of about 5 g/day. Potentially, these supplements could be used to decrease muscle damage or DOMS in athletes. Some groups have noted reductions in DOMS, CK, and inflammatory cytokines (e.g., TNF- $\alpha$ , IL-8) following eccentric contraction induced muscle damage (McFarlin et al., 2016; Nicol et al., 2015). However, similar protective effects of curcumin and reductions in inflammatory makers (e.g., no effect on IL-6, IL-1, IL-10, or CRP) were not seen following endurance exercise (Sciberras et al., 2015). Tart cherry juice has been shown to decrease muscle damage and inflammatory markers following

Manuscript preparation and editing was undertaken by ESR, MPM, and DELM. All authors approved the final version of the paper.

## Downloaded by on 05/29/18, Volume {article.issue.volume}, Article Number {article.issue.issue}

- Downloaded by on 05/29/18, Volume {article.issue.volume}, Article Number {article.issue.issue}

- Fishman, M.P., Lombardo, S.J., & Kharrazi, F.D. (2016). Vitamin D deficiency among professional basketball players. *Orthop J Sports Med*, 4(7), 232596711665574. [PubMed](#) doi:10.1177/2325967116655742
- Fitschen, P.J., Wilson, G.J., Wilson, J.M., & Wilund, K.R. (2013). Efficacy of beta-hydroxy-beta-methylbutyrate supplementation in elderly and clinical populations. *Nutrition*, 29(1), 29–36.
- Fransen, J.C., Zuhl, M., Kerksick, C.M., Cole, N., Altobelli, S., Kuethe, D.O., & Schneider, S. (2015). Impact of creatine on muscle performance and phosphagen stores after immobilization. *European Journal of Applied Physiology*, 115(9), 1877–1886. doi:10.1007/s00421-015-3172-2
- Gentles, J.A., & Phillips, S.M. (2017). Discrepancies in publications related to HMB-FA and ATP supplementation. *Nutrition & Metabolism*, 14, 42. [PubMed](#) doi:10.1186/s12986-017-0201-7
- Ginde, A.A., Mansbach, J.M., & Camargo, C.A., Jr. (2009). Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Archives of Internal Medicine*, 169(4), 384–390. [PubMed](#) doi:10.1001/archinternmed.2008.560
- Gleeson, M., Bishop, N.C., & Struszcak, L. (2016). Effects of Lactobacillus casei Shirota ingestion on common cold infection and herpes virus antibodies in endurance athletes: A placebo-controlled, randomized trial. *European Journal of Applied Physiology*, 116(8), 1555–1563. [PubMed](#) doi:10.1007/s00421-016-3415-x
- Gravina, L., Brown, F.F., Alexander, L., Dick, J., Bell, G., Witard, O.C., & Galloway, S.D. (2017). n-3 fatty acid supplementation during 4 weeks of training leads to improved anaerobic endurance capacity, but not maximal strength, speed, or power in soccer players. *International Journal of Sport Nutrition and Exercise Metabolism*, 27(4), 305–313. [PubMed](#) doi:10.1123/ijnsnem.2016-0325
- Gray, P., Chappell, A., Jenkinson, A.M., Thies, F., & Gray, S.R. (2014). Fish oil supplementation reduces markers of oxidative stress but not muscle soreness after eccentric exercise. *International Journal of Sport Nutrition and Exercise Metabolism*, 24(2), 206–214. [PubMed](#) doi:10.1123/ijnsnem.2013-0081
- Greenhaff, P.L., Bodin, K., Söderlund, K., & Hultman, E. (1994). Effect of oral creatine supplementation on skeletal muscle phosphocreatine re-synthesis. *American Journal of Physiology*, 266(5 Pt 1), E725–E730. [PubMed](#)
- Gualano, B., Rawson, E.S., Candow, D.G., & Chilibeck, P.D. (2016). Creatine supplementation in the aging population: Effects on skeletal muscle, bone and brain. *Amino Acids*, 48(8), 1793–1805. [PubMed](#) doi:10.1007/s00726-016-2239-7
- Gualano, B., Roschel, H., Lancha-Jr, A.H., Brightbill, C.E., & Rawson, E.S. (2012). In sickness and in health: The widespread application of creatine supplementation. *Amino Acids*, 43(2), 519–529. [PubMed](#) doi:10.1007/s00726-011-1132-7
- Halliday, T.M., Peterson, N.J., Thomas, J.J., Kleppinger, K., Hollis, B.W., & Larson-Meyer, D.E. (2011). Vitamin D status relative to diet, lifestyle, injury, and illness in college athletes. *Medicine & Science in Sports & Exercise*, 43(2), 335–343. [PubMed](#) doi:10.1249/MSS.0b013e3181eb9d4d
- Hamilton, B., Grantham, J., Racinais, S., & Chalabi, H. (2010). Vitamin D deficiency is endemic in Middle Eastern sportsmen. *Public Health Nutrition*, 13(10), 1528–1534. [PubMed](#) doi:10.1017/S136898000999320X
- Harris, R.C., Söderlund, K., & Hultman, E. (1992). Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clinical Science (London, England: 1979)*, 83(3), 367–374. doi:10.1042/cs0830367
- Häussinger, D., Roth, E., Lang, F., & Gerok, W. (1993). Cellular hydration state: An important determinant of protein catabolism in health and disease. *Lancet*, 341(8856), 1330–1332. doi:10.1016/0140-6736(93)90828-5
- Haywood, B.A., Black, K.E., Baker, D., McGarvey, J., Healey, P., & Brown, R.C. (2014). Probiotic supplementation reduces the duration and incidence of infections but not severity in elite rugby union players. *Journal of Science and Medicine in Sport*, 17(4), 356–360. [PubMed](#) doi:10.1016/j.jsams.2013.08.004
- He, C.S., Aw Yong, X.H., Walsh, N.P., & Gleeson, M. (2016). Is there an optimal vitamin D status for immunity in athletes and military personnel? *Exercise Immunology Review*, 22, 42–64. [PubMed](#)
- He, C.S., Fraser, W.D., Tang, J., Brown, K., Renwick, S., Rudland-Thomas, J., . . . Gleeson, M. (2016). The effect of 14 weeks of vitamin D3 supplementation on antimicrobial peptides and proteins in athletes. *Journal of Sports Sciences*, 34(1), 67–74. [PubMed](#) doi:10.1080/02640414.2015.1033642
- He, C.S., Handzlik, M., Fraser, W.D., Muhamad, A., Preston, H., Richardson, A., & Gleeson, M. (2013). Influence of vitamin D status on respiratory infection incidence and immune function during 4 months of winter training in endurance sport athletes. *Exercise Immunology Review*, 19, 86–101. [PubMed](#)
- Heaton, L.E., Davis, J.K., Rawson, E.S., Nuccio, R.P., Witard, O.C., Stein, K.W., . . . Baker, L.B. (2017). Selected in season nutritional strategies to enhance recovery for team sport athletes. *Sports Medicine*, 47(11), 2201–2218.
- Hector, A., & Phillips, S.M. (2018). Protein recommendations for weight loss in elite athletes: A focus on body composition and performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 28. doi:10.1123/ijnsnem.2017-0273
- Hespeel, P., Op't Eijnde, B., Van Leemputte, M., Urso, B., Greenhaff, P.L., Labarque, V., . . . Richter, E.A. (2001). Oral creatine supplementation facilitates the rehabilitation of disuse atrophy and alters the expression of muscle myogenic factors in humans. *Journal of Physiology*, 536(Pt 2), 625–633. [PubMed](#) doi:10.1111/j.1469-7793.2001.0625c.xd
- Holick, M.F., Binkley, N.C., Bischoff-Ferrari, H.A., Gordon, C.M., Hanley, D.A., Heaney, R.P., . . . Endocrine, S. (2011). Evaluation, treatment, and prevention of vitamin D deficiency: An Endocrine Society clinical practice guideline. *Journal of Clinical Endocrinology and Metabolism*, 96(7), 1911–1930. [PubMed](#) doi:10.1210/jc.2011-0385
- Hossein-nezhad, A., & Holick, M.F. (2013). Vitamin D for health: A global perspective. *Mayo Clinic Proceedings*, 88(7), 720–755. doi:10.1016/j.mayocp.2013.05.011
- Hultman, E., Söderlund, K., Timmons, J.A., Cederblad, G., & Greenhaff, P.L. (1996). Muscle creatine loading in men. *J Appl Physiol* (1985), 81(1), 232–237. doi:10.1152/jappl.1996.81.1.232
- Jeromson, S., Gallagher, I.J., Galloway, S.D., & Hamilton, D.L. (2015). Omega-3 fatty acids and skeletal muscle health. *Marine Drugs*, 13(11), 6977–7004. [PubMed](#) doi:10.3390/md13116977
- Johnston, A.P., Burke, D.G., MacNeil, L.G., & Candow, D.G. (2009). Effect of creatine supplementation during cast-induced immobilization on the preservation of muscle mass, strength, and endurance. *Journal of Strength & Conditioning Research*, 23(1), 116–120. [PubMed](#) doi:10.1519/JSC.0b013e318181efbcc
- Jouris, K.B., McDaniel, J.L., & Weiss, E.P. (2011). The effect of omega-3 fatty acid supplementation on the inflammatory response to eccentric strength exercise. *Journal of Sports Science & Medicine*, 10(3), 432–438. [PubMed](#)
- Kekkonen, R.A., Vasankari, T.J., Vuorimaa, T., Hahtela, T., Julkunen, I., & Korpela, R. (2007). The effect of probiotics on respiratory infections and gastrointestinal symptoms during training in marathon runners. *International Journal of Sport Nutrition and Exercise Metabolism*, 17(4), 352–363. [PubMed](#) doi:10.1123/ijnsnem.17.4.352

- IJSNEM Vol. 28, No. 2, 2018



- Tipton, K.D. (2015). Nutritional support for exercise-induced injuries. *Sports Medicine*, 45(Suppl 1), 93–104. doi:[10.1007/s40279-015-0398-4](https://doi.org/10.1007/s40279-015-0398-4)
- Turner, C.E., Byblow, W.D., & Gant, N. (2015). Creatine supplementation enhances corticomotor excitability and cognitive performance during oxygen deprivation. *Journal of Neuroscience*, 35(4), 1773–1780. PubMed doi:[10.1523/JNEUROSCI.3113-14.2015](https://doi.org/10.1523/JNEUROSCI.3113-14.2015)
- Vagnozzi, R., Signoretti, S., Floris, R., Marziali, S., Manara, M., Amorini, A.M., . . . Tavazzi, B. (2013). Decrease in N-acetylaspartate following concussion may be coupled to decrease in creatine. *Journal of Head Trauma Rehabilitation*, 28(4), 284–292. PubMed doi:[10.1097/HTR.0b013e3182795045](https://doi.org/10.1097/HTR.0b013e3182795045)
- Volek, J.S., & Rawson, E.S. (2004). Scientific basis and practical aspects of creatine supplementation for athletes. *Nutrition*, 20(7–8), 609–614. doi:[10.1016/j.nut.2004.04.014](https://doi.org/10.1016/j.nut.2004.04.014)
- West, N.P., Pyne, D.B., Cripps, A.W., Hopkins, W.G., Eskesen, D.C., Jairath, A., . . . Fricker, P.A. (2011). Lactobacillus fermentum (PCC (R)) supplementation and gastrointestinal and respiratory-tract illness symptoms: A randomised control trial in athletes. *Nutrition Journal*, 10, 30. PubMed doi:[10.1186/1475-2891-10-30](https://doi.org/10.1186/1475-2891-10-30)
- West, N.P., Pyne, D.B., Peake, J.M., & Cripps, A.W. (2009). Probiotics, immunity and exercise: A review. *Exercise Immunology Review*, 15, 107–126. PubMed
- Wilkinson, D.J., Hossain, T., Hill, D.S., Phillips, B.E., Crossland, H., Williams, J., . . . Atherton, P.J. (2013). Effects of leucine and its metabolite beta-hydroxy-beta-methylbutyrate on human skeletal muscle protein metabolism. *Journal of Physiology*, 591(11), 2911–2923. PubMed doi:[10.1113/jphysiol.2013.253203](https://doi.org/10.1113/jphysiol.2013.253203)
- Willoughby, D.S., & Rosene, J. (2001). Effects of oral creatine and resistance training on myosin heavy chain expression. *Medicine & Science in Sports & Exercise*, 33(10), 1674–1681. PubMed doi:[10.1097/00005768-200110000-00010](https://doi.org/10.1097/00005768-200110000-00010)
- Willoughby, D.S., & Rosene, J.M. (2003). Effects of oral creatine and resistance training on myogenic regulatory factor expression. *Medicine & Science in Sports & Exercise*, 35(6), 923–929. doi:[10.1249/01.MSS.0000069746.05241.F0](https://doi.org/10.1249/01.MSS.0000069746.05241.F0)
- Wilson, J.M., Lowery, R.P., Joy, J.M., Andersen, J.C., Wilson, S.M., Stout, J.R., . . . Rathmacher, J. (2014). The effects of 12 weeks of beta-hydroxy-beta-methylbutyrate free acid supplementation on muscle mass, strength, and power in resistance-trained individuals: A randomized, double-blind, placebo-controlled study. *European Journal of Applied Physiology*, 114(6), 1217–1227. doi:[10.1007/s00421-014-2854-5](https://doi.org/10.1007/s00421-014-2854-5)
- Wilson, J.M., Lowery, R.P., Joy, J.M., Walters, J.A., Baier, S.M., Fuller, J.C., Jr., . . . Rathmacher, J. (2013). Beta-hydroxy-beta-methylbutyrate free acid reduces markers of exercise-induced muscle damage and improves recovery in resistance-trained men. *British Journal of Nutrition*, 110(3), 538–544. doi:[10.1017/S0007114512005387](https://doi.org/10.1017/S0007114512005387)
- Wu, H., Xia, Y., Jiang, J., Du, H., Guo, X., Liu, X., . . . Niu, K. (2015). Effect of beta-hydroxy-beta-methylbutyrate supplementation on muscle loss in older adults: A systematic review and meta-analysis. *Archives of Gerontology and Geriatrics*, 61(2), 168–175. PubMed doi:[10.1016/j.archger.2015.06.020](https://doi.org/10.1016/j.archger.2015.06.020)
- Yquel, R.J., Arsac, L.M., Thiaudiere, E., Canioni, P., & Manier, G. (2002). Effect of creatine supplementation on phosphocreatine resynthesis, inorganic phosphate accumulation and pH during intermittent maximal exercise. *Journal of Sports Sciences*, 20(5), 427–437. PubMed doi:[10.1080/026404102317366681](https://doi.org/10.1080/026404102317366681)
- Zanchi, N.E., Gerlinger-Romero, F., Guimaraes-Ferreira, L., de Siqueira Filho, M.A., Felitti, V., Lira, F.S., . . . Lancha, A.H. Jr. (2011). HMB supplementation: Clinical and athletic performance-related effects and mechanisms of action. *Amino Acids*, 40(4), 1015–1025.